#### RELATION OF WINDS TO TEMPERATURE IN CENTRAL OHIO.

By Howard H. Martin, Observer.

[U. S. Weather Bureau, Columbus, Ohio, Mar. 22, 1920.]

#### SYNOPSIS.

This paper presents by table and graph the existing relation between the direction of the wind and the existing or current temperature, as well as the subsequent 24-hour temperature change. The data are taken from the records of the Columbus (Ohio) office, 1909–1918, inclusive.

It is a fact of common knowledge that some winds are warmer than others, yet it is not generally known that from certain directions, in central Ohio, winds are cooler during the entire year than from others. A glance at figure 1 will reveal this difference, for during all seasons winds from the north and the northwest are accompanied




Fig. 1.—Showing the relation between the prevailing direction of wind at 7 a. m-and 7 p. m. and current temperature, Columbus, Ohio, 1909-1918, inclusive. Warm winds represented by that portion in solid black outside of circle; cold winds by that portion inside. (Scale: 1 inch radially equals 20° F.)

by lower temperatures than from any of the other directions, and also it is evident that winds from the south are warmer than from other directions. The southeast wind may be classified as a warm wind during most of the year, but is cooler than the average of all directions during the summer months. The east wind, too, is much cooler in the summer months, relatively speaking, than during the autumn and winter months.

Table 1 shows the average temperature at the semidaily observation hours, arranged according to the direction of the surface wind at that time. Thus it will be seen again that the north wind for the year is cooler and the south wind warmer than the average. It is to be noted, however, that during the winter months central Ohio experiences the coldest weather with the northeast wind, changing to the northwest in the spring, and to the north in the summer and autumn. During the winter the higher temperatures accompany the southeast wind, changing to the south in spring, to the southwest in summer, and back to the south in the autumn.

There also exists a relation between the direction of the surface wind and the subsequent 24-hour temperature change. That is to say, certain winds may be considered as fair prognostics of colder weather or of warmer weather. Thus, considering figure 2, it will be seen that during the spring months, the southeast, east, northeast, and north directions are followed by warmer weather, the north-




Fig. 2.—Showing the relation between the prevailing direction of wind at 7 a. m. and 7 p. m. and subsequent 24-hour temperature changes, Columbus, Ohio, 1909—1918, in. clusive. Positive changes represented by that portion of wind rose outside of circle; negative changes by that portion on inside. (Scale: 1 inch radially equals 20° F.)

west being practically neutral or without prognostic value, while all other directions presage cooler conditions. During the summer the east wind loses its prognostic value. During the autumn the northwest wind becomes a harbinger of cooler weather, while the east wind again precedes warmer atmosphere. In the winter, it is to be noted, any easterly direction from south to north may be considered as of fair prognostic value indicating increasing temperatures, while all westerly directions indicate lower temperatures.

Table 2 merits careful consideration. It shows the number of times temperature changes, positive and negative, have occurred during the ten years, 1909–1918, inclusive, during the 24 hours following the observation of a given direction of surface wind. Of course, in no case is the prognostic value sufficiently high to warrant implicit

confidence in the wind directions as temperature indicators, but the figures show that there does exist a relation. Especially important is the fact disclosed in Table 2 that of the 32 winter temperature falls equaling or exceeding 30° during the 10-year period in question, 21 occurred following the west and southwest winds. Cold waves during winter months with temperature falls exceeding 20° occurred 123 times, and 85 of these occasions followed a westerly wind. It is also noteworthy that during the winter months 26 of the 123 temperature falls equaling or exceeding 20° followed the southeast wind, normally an indication of higher temperature. This fact was brought about by the passage of rapidly moving Lows, far in excess of normal speed, and the temperature rise as well as the subsequent cold wave occurred within the 24-hour period.

Under ordinary conditions, at Columbus, Ohio, it seems safe to consider the north, northeast, east, and southeast winds as prognostics of warmer weather 24 hours later, except in summer, when the east wind has no prognostic value; and the northwest, west, southwest, and south winds as prognostics of cooler conditions 24 hours later, except in summer, when the northwest wind is usually followed by higher temperatures. However, excepting the case of the east wind in winter, the true prognostic values are low and do not justify much reliance. Their value would be appreciable only when used in conjunction with other indications.

Table 1.—Showing the relation of the 7 a.m. and 7 p.m. direction of wind to the actual air temperature at those hours, and to the average subsequent 24-hour temperature changes, based on the records at Columbus, Ohio, 1909–1918, inclusive.

# TO CURRENT TEMPERATURE.

Direction.	Spring.	Summer.	Autumn.	Winter.	Annual.	
	• F.					
North	44. 5	63.8	46.5	23.0	44.5	
Northeast	46. 4	70.4	52.6	€1.8	47.8	
E <b>ást</b> ,	53.1	70.6	56.8	30.6	52.8	
Southeast	52. 2	68.6	56.2	32. 8	52. 5	
Bouth	58.7	72.5	57.7	32, 5	55.4	
Southwest	57. 6	74.7	55.7	31.6	54.9	
West	48.4	72.1	49.5	26. 1	49.0	
Northwest	41.4	66.3	47.9	23.8	44.8	
Меап, seasonal	50.3	69.8	52.8	28.0	50, 2	

## TO SUBSEQUENT 24-HOUR TEMPERATURE CHANGES.

North Northeast East Southeast Southwest West. Northwest	+2.4 +4.4 +2.6 -4.0 -2.6 -2.0	+2.2 +0.4 +0.1 +1.6 -0.5 -4.4 -0.5 +0.3	+1.3 +4.1 +3.4 +1.7 -2.4 -1.6 -1.8 -1.1	-0.7 +5.3 +7.9 +2.0 +1.1 -3.4 -1.6 -2.0	+1.9 +3.1 +3.9 +2.0 -1.5 -3.0 -1.5 -0.3
Seasonal mean	+0.7	0	0.4	+1.2	+0.6

TABLE 2.—Showing the number of times various wind directions were followed by positive or negative temperature changes, arranged according to the seasons, Columbus, Ohio, period 1909-1918, inclusive.

Direction.	Num-	6 degrees or more.		8 degrees or more.		10 degrees or more.		15 degrees or more.		20 degrees or more.		30 degrees or more.	
	1 640	times noted.	+	_	+	-	+	-	+	_	+	_	+
SPRING.													
North	235 131 159 253 226 214 214 317	61 34 62 131 41 32 30 78	15 13 7 62 97 84 68 68	52 18 33 108 30 27 28 62	9 8 23 48 81 60 47 40	40 12 20 80 19 18 10 53	2 0 44 76 54 25 28	14 4 18 36 13 11 3 20	0 0 30 39 28 14 15	1 0 3 7 4 6 0 3	0 0 7 10 18 4	00010000	0 0 1 2 2 1
SUMMER.													
North Northeast East Southeast South Southwest West Northwest	194 229 320	62 13 20 29 25 17 87 20	28 17 28 17 53 87 27 29	30 5 12 14 19 12 18 13	13 11 23 10 43 59 24 18	17 1 4 6 11 8 10 6	6 3 14 4 17 27 13 9	7 0 0 3 6 2 2 2	1 2 1 8 15 7	1 0 0 1 0 0	0 1 0 1 5 1	000000	0000000
AU <b>TUM</b> N.				1					ļ			Į.	
North Northeast East Southeast South Southwest West Northwest	129 200	41 46 90 75 40 38 24 23	47 21 21 38 88 68 97 31	21 26 72 58 85 28 21 17	31 12 7 35 74 59 74 25	12 19 51 87 25 22 17	26 8 5 30 66 42 53 17	11 32 18 17 13 7	5 1 23 41 31 34 13	0 3 17 7 3 2 3	3 1 0 11 25 16 16	0 2 8 0 1 0 0	0 0 0 0 9 5 3 2
WINTER.											1		
North	123 275 272 280 318	31 65 142 108 90 88 37	44 24 20 72 104 104 129 85	26 28 55 126 82 76 72 26	28 14 15 57 62 97 98 74	22 22 47 105 63 67 62 16	13 10 10 48 50 68 74 62	15 13 33 58 29 35 26 6	7 4 3 36 26 65 46 33	6 6 19 31 14 14 14 0	26 8 32 35 18	1 1 2 6 1 3 0	0 0 3 4 11 10 4

THE VALUE OF HIGH-LEVEL METEOROLOGICAL DATA IN FORECASTING CHANGES OF TEMPERATURE—A CONTRIBUTION TO THE METEOROLOGY OF MOUNT ROSE, NEV.<sup>1</sup>

By S. P. Fergusson.

### [Author's summary.]

The general relation or connection between the conditions recorded at the summit and base stations of Mount Rose appears to be practically the same as that found to exist between the summit and base stations of mountains in other parts of the world.

Of the decided falls of temperature or cold waves occurring on the summit during four years of observa-

<sup>&</sup>lt;sup>1</sup> Reprinted from Technical Bulletin No. 83, University of Nevada Agricultural Experiment Station, Reno, Nev., 1915, pp. 29 and 30.